







Name: \_\_\_\_\_

Period: \_\_\_\_\_

**Advanced Forces Practice**


1. Match the following equations with the correct masses. These equations could be in either the x or y-direction. There is one duplicate.

- A. \_\_\_\_\_  $T = ma$
- B. \_\_\_\_\_  $T - T - F_W = ma$
- C. \_\_\_\_\_  $F_N - F - F_W = ma$
- D. \_\_\_\_\_  $T - T \cos \theta = 0$
- E. \_\_\_\_\_  $T - F_f = ma$
- F. \_\_\_\_\_  $F_W \sin \theta = ma$
- G. \_\_\_\_\_  $F_N + F - F_W = ma$
- H. \_\_\_\_\_  $F_N - F_W = mv^2/r$
- I. \_\_\_\_\_  $T \sin \theta - F_W = 0$
- J. \_\_\_\_\_  $F_N + F - F - F_W = ma$
- K. \_\_\_\_\_  $T - F_f = 0$
- L. \_\_\_\_\_  $T - T - F_f = ma$
- M. \_\_\_\_\_  $F_W = ma$
- N. \_\_\_\_\_  $T - F_W = ma$
- O. \_\_\_\_\_  $F_N + F_W = mv^2/r$
- P. \_\_\_\_\_  $F_N - F_W = ma$






- Q. \_\_\_\_\_  $F - T = ma$
- R. \_\_\_\_\_  $F_N = mv^2/r$
- S. \_\_\_\_\_  $F_N - mg \cos \theta = ma$
- T. \_\_\_\_\_  $F_W \sin \theta - T = ma$

2. A 15 kg mass has a 45 N force pulling on it at an angle of  $48^\circ$  above the horizon. The mass is on a surface that has the following coefficients of friction:  $\mu_s = 0.34$  and  $\mu_k = 0.16$ .
- A. Decide if it will move, if it starts at rest.
  - B. Calculate the acceleration if it is already moving.




3. A. Calculate how heavy the 65 kg lady in the elevator feels.
- B. What would a scale (reading weight) read that is under her feet?
- C. What would a scale read if the elevator's cable was cut?
4. Slim Jim's dog "Bim" is pulling 15N on a 5.2 kg mass at a constant velocity of 2 m/s. There is friction between the mass and the floor.
- A. Draw and label all of the forces acting on the mass.
  - B. What is the acceleration of the object?
  - C. Calculate the force of friction on the mass.
  - D. Calculate the coefficient of friction of the floor.





5. A 25 kg object has a velocity of  $-12\text{ m/s}$  and has an acceleration of  $+2\text{ m/s}^2$ .
- Is the object moving to the left, to the right, or at rest?
  - Is the object speeding up or slowing down?
  - Are the forces balanced or unbalanced?
  - How do you know?
  - Which force is greater:  $F_1$  or  $F_2$ ?
  - Calculate the net force acting on the object.

G. Calculate the magnitude of force 2.



7. Slim Jim has a rope attached to an 40 kg box.
- If the box is not moving or at constant speed, what is its acceleration?
  - What is the tension in the rope?
  - If Slim Jim pulls the object up with an acceleration of  $2.5\text{ m/s}^2$ , find the tension in the rope.

8. Find the acceleration for each of the 5 kg masses below. On the right there is only one mass and Slim Jim pulls down with 200 N.

